



[Go to Product page](#)

Datasheet for ABIN1043792

anti-Interleukin 17a antibody (Biotin)

1 Image

1 Publication

Overview

Quantity:	100 µg
Target:	Interleukin 17a (IL17A)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Interleukin 17a antibody is conjugated to Biotin
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	Anti-IL-17A (MOUSE) Monoclonal Antibody was produced in mouse by repeated immunizations with mature full length recombinant human IL-17A produced in E.coli followed by hybridoma development. Immunogen Type: RecombinantProtein
Clone:	4H1524-1
Isotype:	IgG2b
Specificity:	This product was purified from concentrated tissue culture supernate by Protein G chromatography followed by extensive dialysis against the buffer stated above. This antibody is specific for human IL-17A protein. A BLAST analysis was used to suggest cross-reactivity with IL-17A from human sources based on 100% homology with the immunizing sequence. Cross-reactivity with IL-17A from other sources has not been determined.
Characteristics:	IL17-A (also known as Interleukin-17) is a proinflammatory cytokine produced by activated T

Product Details

cells. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. IL17-A is the founding member of a group of cytokines called the IL17 family. IL17-A was originally identified as a transcript from a rodent T-cell hybridoma. To elicit its functions, IL17 binds to a type I cell surface receptor called IL17R of which there are at least three variants IL17RA, IL17RB, and IL17RC. Anti-IL-17A antibody is ideal for investigators involved in Immunology research.

Purification: purified

Target Details

Target: Interleukin 17a (IL17A)

Alternative Name: IL-17A ([IL17A Products](#))

Background: IL17-A (also known as Interleukin-17) is a proinflammatory cytokine produced by activated T cells. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. IL17-A is the founding member of a group of cytokines called the IL17 family. IL17-A was originally identified as a transcript from a rodent T-cell hybridoma. To elicit its functions, IL17 binds to a type I cell surface receptor called IL17R of which there are at least three variants IL17RA, IL17RB, and IL17RC. Anti-IL-17A antibody is ideal for investigators involved in Immunology research.

Synonyms: Interleukin-17, IL-17, IL-17A, Interleukin-17A, Cytotoxic T-Lymphocyte-associated Antigen 8, CTLA8.

Gene ID: 3605

NCBI Accession: [NP_002181](#)

UniProt: [Q16552](#)

Application Details

Application Notes: This purified antibody has been tested for use in FLOW and Western Blot. Specific conditions for reactivity should be optimized by the end user.

Application Details

Comment: Gene Name: IL17A

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Reconstitution Buffer: Restore with deionized water (or equivalent), Reconstitution Volume: 100 μ L

Concentration: 1.0 mg/mL

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Preservative: Sodium azide

Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C

Storage Comment: Store vial at 4° C before opening. DO NOT FREEZE. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. Freezing alkaline phosphatase conjugates will result in a substantial loss of enzymatic activity. Expiration date is one (1) year from date of opening.

Expiry Date: 12 months

Publications

Product cited in: Baek, Eling: "Changes in gene expression contribute to cancer prevention by COX inhibitors." in: **Progress in lipid research**, Vol. 45, Issue 1, pp. 1-16, (2006) ([PubMed](#)).



Western Blotting

Image 1. Western Blot of Mouse Anti-IL-17A antibody Lane 1: human full length recombinant IL-17A protein Lane 2: mouse full length recombinant IL-17A protein Lane 3: rat full length recombinant IL-17A protein Load: 20 ng/lane Primary antibody: Anti-IL-17A antibody at 1ug/mL for overnight at 4°C